

Publication List Franca Fraternali

- [1] Autore F, Bergeron JRC, Malim MH, Fraternali F and Huthoff H. Rationalisation of the differences between APOBEC3G structures from crystallography and NMR studies by molecular dynamics simulations. *PLoS ONE* **5** e11515 (2010).
- [2] Autore F, Pagano B, Fornili A, Rittinger K and Fraternali F. In Silico Phosphorylation of the Autoinhibited Form of p47(phox): Insights into the Mechanism of Activation. *Biophysical Journal* **99**(11) 3716–3725 (2010).
- [3] Chakroun N, Prigent S, Dreiss CA, Noinville S, Chapuis C, Fraternali F and Rezaei H. The oligomerization properties of prion protein are restricted to the H2H3 domain. *FASEB J.* **24** 3222–3231 (2010).
- [4] Fernandes LP, Annibale A, Kleinjung J, Coolen ACC and Fraternali F. Protein networks reveal detection bias and species consistency when analysed by information-theoretic methods. *PLoS ONE* **5** e12083 (2010).
- [5] Merabet A, Houllberghs H, Maclagan K, Akanho E, Bui TTT, Pagano B, Drake AF, Fraternali F and Nikolova PV. Mutants of the tumour suppressor p53 L1 loop as second-site suppressors for restoring DNA binding to oncogenic p53 mutations: structural and biochemical insights. *Biochem J* **427** 225–236 (2010).
- [6] Vaz F, Hanenberg H, Schuster B, Barker K, Wiek C, Erven V, Neveling K, Endt D, Kesterton I, Autore F, Fraternali F, Freund M *et al.* Mutation of the RAD51C gene in a Fanconi anemia-like disorder. *Nat. Genet.* **42** 406–409 (2010).
- [7] Annibale A, Coolen ACC, Fernandes LP, Fraternali F and Kleinjung J. Tailored graph ensembles as proxies or null models for real networks I: tools for quantifying structure. *Journal of Physics A: Math. Theor.* **42** 485001 (25 pp) (2009).
- [8] Huthoff H, Autore F, Gallois-Montbrun S, Fraternali F and Malim MH. RNA-dependent oligomerization of APOBEC3G is required for restriction of HIV-1. *PLoS Pathog.* **5** e1000330 (2009).
- [9] Nicastro G, Masino L, Esposito V, Menon R, Simone AD, Fraternali F and Pastore A. Josephin domain of ataxin-3 contains two distinct ubiquitin-binding sites. *Biopolymers* **91** 1203–1214 (2009).
- [10] Coluzza I, De Simone A, Fraternali F and Frenkel D. Multi-scale simulations provide supporting evidence for the hypothesis of intramolecular protein translocation in GroEL/GroES complexes. *PLoS Comput. Biol.* **4** e1000006 (2008).
- [11] Festa G, Autore F, Fraternali F, Giardina P and Sannia G. Development of new laccases by directed evolution: functional and computational analyses. *Proteins* **72** 25–34 (2008).
- [12] Pagano B, Mattia CA, Cavallo L, Uesugi S, Giancola C and Fraternali F. Stability and cations coordination of DNA and RNA 14-mer G-quadruplexes: a multiscale computational approach. *J. Phys. Chem. B* **112** 12115–12123 (2008).
- [13] Patel S, Bui TTT, Drake AF, Fraternali F and Nikolova PV. The p73 DNA binding domain displays enhanced stability relative to its homologue, the tumor suppressor p53, and exhibits cooperative DNA binding. *Biochemistry* **47** 3235–3244 (2008).
- [14] Patel S, George R, Autore F, Fraternali F, Ladbury JE and V NP. Molecular interactions of ASPP1 and ASPP2 with the p53 protein family and the apoptotic promoters PUMA and Bax. *Nucleic Acids Res.* **36** 5139–5151 (2008).

- [15] Rabello S, Coolen ACC, Perez-Vicente CJ and Fraternali F. A solvable model of the genesis of amino-acid sequences via coupled dynamics of folding and slow genetic variation. *J. Phys. A: Math. Theor.* **41** 285004 (50pp) (2008).
- [16] Virno A, Mayol L, Ramos A, Fraternali F, Pagano B and Randazzo A. Structural insight into the hTERT intron 6 sequence d(GGGGTGAAAGGGG) from $^1\text{H-NMR}$ study. *Nucleosides Nucleotides Nucleic Acids* **26** 1133–1137 (2008).
- [17] Autore F, Melchiorre S, Kleinjung J, Morgan WD and Fraternali F. Interaction of malaria parasite-inhibitory antibodies with the merozoite surface protein MSP1₁₉ by computational docking. *Proteins* **66** 513–527 (2007).
- [18] Azzu V, Yadin D, Patel H, Fraternali F, Chantler PD and Molloy JE. Calcium regulates scallop muscle by changing myosin flexibility. *Eur. Biophys. J.* **35** 302–312 (2007).
- [19] De Simone A, Spadaccini R, Temussi PA and Fraternali F. Toward the understanding of MNEI sweetness from hydration map surfaces. *Biophys. J.* **90** 3052–3061 (2007).
- [20] Pandini A, Bonati L, Fraternali F and Kleinjung J. MinSet: a general approach to derive maximally representative database subsets by using fragment dictionaries and its application to the SCOP database. *Bioinformatics* **23** 515–516 (2007).
- [21] De Simone A, Dodson GG, Fraternali F and Zagari A. Water molecules as structural determinants among prions of low sequence identity. *FEBS Lett.* **580** 2488–2494 (2006).
- [22] Mollica L, Fraternali F and Musco G. Interactions of the C2 domain of human factor V with a model membrane. *Proteins* **64** 363–375 (2006).
- [23] Taylor WR, Lin K, Klose D, Fraternali F and Jonassen I. Dynamic domain threading. *Proteins* **64** 601–614 (2006).
- [24] De Simone A, Dodson GG, Verma CS, Zagari A and Fraternali F. Prion and water: tight and dynamical hydration sites have a key role in structural stability. *Proc. Natl. Acad. Sci. USA* **102** 7535–7540 (2005).
- [25] Kleinjung J and Fraternali F. Molecular simulations in structure predictions. In Jorde LB, editor, *Encyclopedia of Genetics, Genomics, Proteomics and bioinformatics*, 28.14. John Wiley and Sons, Ltd (2005).
- [26] Kleinjung J and Fraternali F. POPSCOMP: an automated interaction analysis of biomolecular complexes. *Nucleic Acids Res.* **33** W342–W346 (2005).
- [27] Vaccaro L, Cross KJ, Kleinjung J, Straus SK, Thomas DJ, Wharton SA, Skehel JJ and Fraternali F. Plasticity of influenza haemagglutinin fusion peptides and their interaction with lipid bilayers. *Biophys. J.* **88** 25–36 (2005).
- [28] Vaccaro L, Cross KJ, Wharton SA, Skehel JJ and Fraternali F. Aspects of the fusogenic activity of influenza haemagglutinin peptides by molecular dynamics simulations. In Fischer WB, editor, *Viral membrane proteins: structure, function, and drug design*, 65–75. Kluwer Academic / Plenum Publisher (2005).
- [29] Cavallo L, Moore MH, Corrie JET and Fraternali F. Quantum mechanics calculations on rhodamine dyes require inclusion of solvent water for accurate representation of the structure. *J. Phys. Chem. A* **108** 7744–7751 (2004).
- [30] Eberini I, Baptista AM, Gianazza E, Fraternali F and Beringhelli T. Reorganization in apo- and holo- β -lactoglobulin upon protonation of Glu89: molecular dynamics and pKa calculations. *Proteins* **54** 744–758 (2004).

- [31] Mazzini S, Bellucci MC, Dallavalle S, Fraternali F and Mondelli R. Mode of binding of camptothecins to double helix oligonucleotides. *Org. Biomol. Chem.* **2** 505–513 (2004).
- [32] Cavallo L, Kleinjung J and Fraternali F. POPS: A fast algorithm for solvent accessible surface areas at atomic and residue level. *Nucleic Acids Res.* **31** 3364–3366 (2003).
- [33] Fraternali F, Cavallo L and Musco G. Effects of pathological mutations on the stability of a conserved amino acid triad in retinoschisin. *FEBS Lett.* **544** 21–26 (2003).
- [34] Kleinjung J, Fraternali F, Martin SR and Bayley PM. Thermal unfolding simulations of apo-calmodulin using leap-dynamics. *Proteins* **50** 648–656 (2003).
- [35] Mombelli E, Morris R, Taylor W and Fraternali F. Hydrogen-bonding propensities of sphingomyelin in solution and in a bilayer assembly: a molecular dynamics study. *Biophys J.* **84** 1507–1517 (2003).
- [36] Fraternali F and Cavallo L. Parameter optimized surfaces (POPS): analysis of key interactions and conformational changes in the ribosome. *Nucleic Acids Res.* **30** 2950–2960 (2002).
- [37] Amodeo P, Fraternali F, Lesk AM and Pastore A. Modularity and homology: modelling of the titin type I modules and their interfaces. *J. Mol. Biol.* **311** 283–296 (2001).
- [38] Falcigno L, Fraternali F, Manduca DM, D’Auria G, Simonetti M, Di Bello C and Paolillo L. A conformational study in solution of pro-somatostatin fragments by NMR and computational methods. *J. Pept. Sci.* **4** 305–318 (2000).
- [39] Fraternali F, Amodeo P, Musco G, Nilges M and Pastore A. Exploring protein interiors: the role of a buried histidine in the KH module fold. *Proteins* **34** 484–496 (2000).
- [40] Fraternali F, Do QT, Doan BT, Atkinson RA, Palmas P, Sklenar V, Safar P, Wildgoose P, Strop P and Saudek V. Mapping the active site of factor Xa by selective inhibitors: an NMR and MD study. *Proteins* **30** 264–274 (2000).
- [41] Kleinjung J, Bayley P and Fraternali F. Leap-dynamics: efficient sampling of conformational space of proteins and peptides in solution. *FEBS Lett.* **470** 257–262 (2000).
- [42] Adinolfi S, Bagni C, Castiglione Morelli MA, Fraternali F, Musco G and Pastore A. Novel RNA-binding motif: the KH module. *Biopolymers* **51** 153–164 (1999).
- [43] De Marino S, Morelli MA, Fraternali F, Tamborini E, Musco G, Vrtala S, Dolecek C, Arosio P, Valenta R and Pastore A. An immunoglobulin-like fold in a major plant allergen: the solution structure of Phl p 2 from timothy grass pollen. *Structure* **7** 943–952 (1999).
- [44] Fraternali F, Amodeo P, Musco G, Nilges M and Pastore A. Exploring protein interiors: the role of a buried histidine in the KH module fold. *Proteins: Struct. Func. Gen.* **34** 484–496 (1999).
- [45] Fraternali F, Anselmi C and Temussi PA. Neurologically active plant compounds and peptide hormones: a chirality connection. *FEBS Lett.* **448** 217–220 (1999).
- [46] Fraternali F and Pastore A. Modularity and homology: modelling of the type II module family from titin. *J. Mol. Biol.* **290** 581–593 (1999).

- [47] Cavallo L and Fraternali F. A molecular dynamics study of the first five generations of poly(propyleneimine) dendrimers modified with N-t-Boc-L-phenylalanine. *Chem. Eur. J.* **4** 927–934 (1998).
- [48] Doan BT, Fraternali F, Do QT, Atkinson RA, Palmas P, Sklenar V, Wildgoose P, Strop P and Saudek V. NMR and MD study of the active site of factor Xa by selective inhibitors. *J. Chem. Phys.* **95** 443–446 (1998).
- [49] Crescenzi O, Fraternali F, Picone D, Tancredi T, Balboni G, Guerrini R, Lazarus LH, Salvadori S and Temussi PA. Design and solution structure of a partially rigid opioid antagonist lacking the basic center - models of antagonism. *Eur. J. Biochem.* **247** 66–73 (1997).
- [50] Fraternali F and Wipff G. Binding of NR_4^+ ammonium ions by a bridged calix[6]arene in chloroform. An MD investigation. *Anales de Chimica Int. Ed.* **93** 376–384 (1997).
- [51] Fraternali F and Wipff G. Interaction of the t-butyl-calix[4]arene anion with ammonium cations in acetonitrile: host-guest or exo counterions? A molecular dynamics investigation. *J. Incl. Phenom. Mol. Recog.* **28** 63–78 (1997).
- [52] Fraternali F and Wipff G. A molecular dynamics study of complexation and stability of luminescent cations encapsulating ligands with bipyridine units. *J. Phys. Org. Chem.* **10** 292–304 (1997).
- [53] Musco G, Kharrat A, Stier G, Fraternali F, Gibson TJ, Nilges M and Pastore A. The solution structure of the first KH domain of FMR1, the protein responsible for the fragile X syndrome. *Nat. Struct. Biol.* **4** 712–716 (1997).
- [54] Ulrich G, Ziessel R, Manet I, Guardigli M, Sabbatini N, Fraternali F and Wipff G. Calix[4]arene in podants and barrelands incorporating 2-2'-bipyridinemoieties and their lanthanide complexes: luminescence properties. *Chem. Eur. J.* **3** 1815–1822 (1997).
- [55] Fraternali F and Van Gunsteren WF. An efficient mean solvation force model for use in molecular dynamics simulations of proteins in aqueous solution. *J. Mol. Biol.* **256** 939–948 (1996).
- [56] Benedetti E, Carlomagno T, Fraternali F, Hamada Y, Hayashi K, Paolillo L and Shioiri T. Conformational analysis of dolastatin 10: an NMR and theoretical approach. *Biopolymers* **36** 525–538 (1995).
- [57] Fraternali F and Van Gunsteren WF. Conformational transitions of a dipeptide in water: effects of imposed pathways using umbrella sampling techniques. *Biopolymers* **34** 347–355 (1994).
- [58] Paolillo L, D'Auria G, Falcigno L, Fraternali F, Saviano M, Simonetti M and Di Bello C. Conformational studies on peptide fragments related to oxytocin and somatostatin precursors. *Curr. Top. in Pep. Prot. Res.* **1** 193–222 (1994).
- [59] Van Gunsteren WF, Beutler TC, Fraternali F, King PM, Mark AE and Smith PE. Computation of free energy in practice: choice of approximations and accuracy limiting factors. In Van Gunsteren WF, Weiner PK and Wilkinson AJ, editors, *Computer simulation of biomolecular systems, theoretical and experimental applications*, volume 2, 315–347. Escom Science Publishers BV (1993).
- [60] Pavone V, Lombardi A, Natri F, Saviano M, Di Blasio B, Fraternali F, Pedone A and Yamada T. First observation of a helical peptide containing chiral α -monosubstituted residues without a preferred screw sense. *J. Chem. Soc. - Perkin Trans. II* 971–977 (1992).

- [61] Crisma M, Valle G, Bonora GM, Toniolo C, Lelj F, Barone V, Fraternali F, Hardy PM and Maia HL. Preferred conformation of peptides from $C^{\alpha,\alpha}$ -symmetrically disubstituted glycines: aromatic residues. *Biopolymers* **31** 637–641 (1991).
- [62] Toniolo C, Crisma M, Valle G, Bonora GM, Lelj F, Barone V, Fraternali F and Hardy HLS P M Maia. Conformational analysis of peptides from $C^{\alpha,\alpha}$ -symmetrically disubstituted aromatic α -amino acid residues. In Giralt E and Andreu D, editors, *Peptides*, 458–459. Escom Science Publishers BV (1991).
- [63] Amodeo P, Barone V and Fraternali F. A microscopic approach to the structure and thermodynamic properties of peptides and proteins. *Thermochimica Acta* **162** 141–154 (1990).
- [64] Barone V, Fraternali F and Cristinziano PL. Sensitivity of peptide conformation to methods and geometrical parameters. A comparative *ab initio* and molecular mechanics study of oligomers of α -aminoisobutyric acid. *Macromolecules* **23** 2038–2044 (1990).
- [65] Crisma M, Valle G, Bonora GM, De Menego E, Toniolo C, Lelj F, Barone V and Fraternali F. Linear oligopeptides 223. Structural versatility of peptides from $C^{\alpha,\alpha}$ -disubstituted glycines. Preferred conformation of the $C^{\alpha,\alpha}$ -diphenylglycine residue. *Biopolymers* **30** 1–11 (1990).
- [66] Fraternali F. Restrained and unrestrained molecular dynamics simulations in the NVT ensemble of alamethicin. *Biopolymers* **30** 1083–1099 (1990).
- [67] Valle G, Crisma M, Bonora GM, Toniolo C, Lelj F, Barone V, Fraternali F, Hardy PM and Langran-Goldsmith A. Structural versatility of peptides from $C^{\alpha,\alpha}$ -disubstituted glycines. Preferred conformation of the $C^{\alpha,\alpha}$ -dibenzylglycine residue. *J. Chem. Soc. - Perkin Trans. II* 1481–1487 (1990).
- [68] Zanotti G, Paolillo L, Barbato G, Fraternali F and D’Alagni M. Cation binding to heterodetic peptidic systems: CD and NMR structural studies. In Giralt E and Andreu D, editors, *Peptides*, 499–500. Escom Science Publishers BV (1990).
- [69] Benedetti E, Di Blasio B, Pavone V, Pedone C, Santini A, Barone V, Fraternali F, Lelj F, Bavoso A and Crisma M. Structural versatility of peptides containing $C^{\alpha,\alpha}$ -dialkylated glycines. An X-ray diffraction study of six 1-aminocyclopropane-1-carboxylic acid rich peptides. *Int. J. Biol. Macromol.* **11** 353–360 (1989).
- [70] Crisma M, Bonora GM, Toniolo C, Barone V, Benedetti E, Di Blasio B, Pavone V, Pedone C, Santini A and Fraternali F. Structural versatility of peptides containing $C^{\alpha,\alpha}$ -dialkylated glycines: conformational energy computations, IR absorption and $^1\text{H-NMR}$ analysis of 1-aminocyclopropane-1-carboxylic acid homopeptides. *Int. J. Biol. Macromol.* **11** 345–352 (1989).
- [71] Barone V, Fraternali F, Cristinziano PL, Lelj F and Rosa A. Conformational behavior of $C^{\alpha,\alpha}$ -dialkylated peptides: *ab initio* and empirical results for cyclopropylglycine. *Biopolymers* **27** 1673–1685 (1988).
- [72] Santini A, Barone V, Bavoso A, Benedetti E, Di Blasio B, Fraternali F, Lelj F, Pavone V, Pedone C, Crisma M, Bonora GM and Toniolo C. Structural versatility of peptides from $C^{\alpha,\alpha}$ -dialkylated glycines: a conformational energy calculation and X-ray diffraction study of homopeptides from 1-aminocyclopentane-1-carboxylic acid. *Int. J. Biol. Macromol.* **10** 292–299 (1988).
- [73] Toniolo C, Valle G, Bonora GM, Lelj F, Barone V, Fraternali F, Callet G, Wagnon J and Nisato D. Conformational preferences and self-association modes of two diastereomeric statine derivatives. *Int. J. Pept. Protein. Res.* **30** 583–595 (1987).